

THAT WHICH IS CLAIMED IS:

1. A method of making a metal complex, comprising the steps of:

(a) acylating a dipyrromethane or a 1-monoacyldipyrromethane to form a mixed reaction product comprising a 1,9-diacyldipyrromethane;

5 (b) combining said mixed reaction product with a compound of the formula R_2MX_2 in the presence of a base, where R is alkyl or aryl, M is Sn, Si, Ge, or Pb, and X is halo, OAc, acac or OTf, to form a metal complex of the formula DMR_2 in said mixed reaction product, wherein D is a 1,9-diacyldipyrromethane; and then

(c) separating said metal complex from said mixed reaction product.

10

2. The method of claim 1, wherein said acylating step (a) is carried out by reacting said dipyrromethane or 1-monoacyldipyrromethane with a compound of the formula R^3COX , where R^3 is alkyl or aryl and X is halo, to form said mixed reaction product comprising a 1,9-diacyldipyrromethane acylated at the 1 and 9 positions with
15 R^3CO- .

15

3. The method of claim 1, wherein said acylating step (a) is carried out by reacting said dipyrromethane or 1-monoacyldipyrromethane with an acid chloride and a Grignard reagent to form said mixed reaction product comprising a 1,9-diacyldipyrromethane.
20

20

4. The method of claim 1, wherein said acylating step (a) is carried out by reacting said dipyrromethane or 1-monoacyldipyrromethane with an active ester to form said mixed reaction product comprising a 1,9-diacyldipyrromethane.

25

5. The method of claim 1, wherein said acylating step (a) is carried out by reacting said dipyrromethane or 1-monoacyldipyrromethane with a Vilsmeier reagent to form said mixed reaction product comprising a 1,9-diacyldipyrromethane.

30

6. The method of claim 1, wherein said base is selected from the group consisting of triethylamine, tributylamine, *N,N*-diisopropylamine, DBU, DBN, and 2,6-di-*tert*-butylpyridine.

7. The method of claim 1, wherein M is Sn.

8. The method of claim 1, wherein said acylating step (a) is carried out with a dipyrromethane which is thereby acylated at the 1 and 9 position to produce said 1,9-
5 diacyldipyrromethane.

9. The method of claim 1, wherein said acylating step (a) is carried out with a 1-monoacyldipyrromethane which is acylated at the 9 position to produce said 1,9-
10 diacyldipyrromethane.

10. The method of claim 1, wherein said compound of the formula R_2MX_2 is immobilized on a solid support.

11. The method of claim 1, further comprising the step of:
15 (d) treating said metal complex with an acid to produce a 1,9-
diacyldipyrromethane.

12. The method of claim 11, wherein said acid is selected from the group consisting of trifluoroacetic acid, trichloroacetic acid, acetic acid, HCl, *p*-toluene
20 sulfonic acid.

13. The method of claim 1, further comprising the steps of:
(d) reducing said metal complex with a reducing agent to form a diol from said
1,9-diacyldipyrromethane; and then
25 (e) condensing said diol with a dipyrromethane to form a porphyrin ring
compound therefrom.

14. The method of claim 13, wherein said reducing agent is $NaBH_4$.

30 15. The method of claim 1, wherein said dipyrromethane is substituted at the 5
position with a substituent selected from the group consisting of H, alkyl, and aryl.

16. The method of claim 1, wherein said dipyrromethane is substituted at the 5 position with a substituent selected from the group consisting of dipyrromethane, porphyrin, dipyrin, and diacyldipyrromethane.

5 17. A method of making a metal complex, comprising:

 reacting a 1,9-diacyldipyrromethane with a compound of the formula R_2MX_2 in the presence of a base, where R is alkyl or aryl, M is Sn, Si, Ge or Pb, and X is halo, OAc, acac or OTf, to form a metal complex of the general formula DMR_2 , wherein D is said 1,9-diacyldipyrromethane.

10

 18. The method of claim 17, wherein said base is selected from the group consisting of triethylamine, tributylamine, *N,N*-diisopropylamine, DBU, DBN, and 2,6,-di-*tert*-butylpyridine.

15 19. The method of claim 17, wherein M is Sn.

 20. A metal complex of the general formula DMR_2 , wherein D is a 1,9-diacyldipyrromethane, M is Sn, Si, Ge, or Pb, and R is alkyl or aryl.

20 21. The metal complex of claim 20, wherein M is Sn.

 22. The metal complex of claim 20, wherein said dipyrromethane is substituted at the 5 position with a substituent selected from the group consisting of H, alkyl, and aryl.

25

 23. The metal complex of claim 20, wherein said dipyrromethane is substituted at the 5 position with a substituent selected from the group consisting of dipyrromethane, porphyrin, dipyrin, and diacyldipyrromethane.

30